

Ewing Demonstration Center

Maturity Group IV

2004 Roundup Ready

(200 Series) Soybean Plot



UNIVERSITY OF ILLINOIS
EXTENSION

Harvested area/variety:

Date planted: May 11, 2004
Planting method: No till drill, 18 rows @ 10 inches
Soil conditions at planting: Good
Date herbicides applied:
Burndown:
Pre-emerge: April 30, 2004
Roundup Weathermax 22 oz/ac
Prowl H₂O 32 oz/ac
2, 4 - D LV 8 oz/ac

Post-emerge: June 2, 2004
Roundup Weathermax 22 oz/ac
June 27, 2004
Roundup Weathermax 22 oz/ac

Date harvested: October 9, 2004
Grain moisture: Adjusted to 13%

Seed Source	Variety	Harvest Moisture %	Bushel/ Acre
buffer			
Adler		398 RRN	15.1 61.1
DeKalb	DKB	46-51 RR	14.8 54.1
Bergmann-Taylor	BT	434 CR	14.5 61.3
Baker Seeds		4425 CRR	14.2 64.8
Trisler	T	4838 RRCN	14.2 64.0
Hoffman Seed	H	3441 RRCN	14.3 58.5
Stine		4542-4 RR	14.3 45.1
Steyer		4410 RRSCN	14.1 66.1
FS	HS	4536 RR	13.9 50.5
Asgrow	AG	4801 RR	14.1 70.0
NK	S	49-Q9 RR	13.9 68.6
M & D Seed		9440 RR	14.0 59.1
Hoffman Seed	H	3459 RRCR	13.8 57.6
Stine		4842-4 RR	13.9 55.6
Baker Seeds		4825 CRR	13.7 58.9
Steyer		4000 RRSCN	14.0 53.1
Bergmann-Taylor	BT	484 CR	13.7 45.9
FS	HS	4736 RR	13.7 48.6
Trisler	T	4227 RRCN	13.7 51.8
M & D Seed		9470 RR	13.8 40.3
buffer			

Plot Average (200 Series)

56.8 bu/ac